

Modular Invariants of $N = 2$ Supersymmetric $SU(1, 1)$ Models

Katri Huitu^{*†}

Department of Physics
University of Southern California
University Park
Los Angeles, CA 90089-0484

We study the modular invariance of $N = 2$ superconformal $SU(1, 1)$ models. By decomposing the characters of Kazama-Suzuki model $SU(3)/(SU(2) \times U(1))$ into an infinite sum of the characters of $(SU(1, 1)/U(1)) \times U(1)$ we construct modular invariant partition functions of $(SU(1, 1)/U(1)) \times U(1)$.

^{*} Supported in part by a grant from the Emil Aaltonen Foundation

[†] Address after June 1: Univ. of Helsinki, Research Institute for High Energy Physics, Siltaavuorenpenger 20 C, 00170 Helsinki, Finland

1. Introduction

$N = 2$ superconformal theories with $c > 3$ are phenomenologically attractive as potentially realistic internal sectors of string compactifications. Two types of models of this kind are known: the Kazama-Suzuki models based on coset construction of affine Lie algebras [1] and non-compact models [2], [3], [4]. In [2] it was shown that all unitary $N = 2$ models with $c > 3$ can be written as

$$\frac{SU(1,1)}{U(1)} \times U(1). \quad (1.1)$$

A stringent restriction for conformal theories comes from modular invariance. The modular properties of characters of Kazama-Suzuki models without fixed points of field identifications are well known [5]. On the other hand the modular invariance remains largely an open question for non-compact models. Some attempts to answer these questions for strings on $SU(1,1)$ manifold have been investigated in [6]. It was suggested that the characters of the discrete series $D^\pm$ contain new sectors, which correspond to isomorphic algebras with integer twists. It was argued that then the left-right symmetric product of the $D^+ \oplus D^-$ characters is a modular invariant.

Another way of finding modular invariant partition functions for the $SU(1,1)$ models is to express a known modular invariant partition function in terms of non-compact characters. For instance one could use this approach to determine the field content of the parafermionic coset model $SU(1,1)/U(1)$ at level $9/4$ or central charge $c = 26$ [4]. This model is of great interest, since it was identified by Witten [7] as a two-dimensional black hole*. The coset structure (1.1) indicates that the $N = 2$ partition function of $SU(1,1)$ can be expressed in terms of the parafermionic partition function and the bosonic partition function.

As an example of these ideas we study the $N = 2$ superconformal $SU(1,1)$ models. In particular, we show that the characters of the Kazama-Suzuki model $SU(3)/(SU(2) \times U(1))$ can be expressed as an infinite sum of characters of $SU(1,1)$ model. Using the relationship between these two theories it is clear that a modular invariant combination of the $SU(1,1)$ characters is given by the diagonal $SU(3)/(SU(2) \times U(1))$ partition function.

* The string theories in curved space-time were also discussed in [4], and it was proposed that $SO(d-1,2)_{-k}/SO(d-1,1)_{-k}$ describe strings propagating in $(d-1)+1$ dimensions.

2. $N = 2$ Characters of $SU(1, 1)$

It is well known that minimal $N = 2$ models with $c < 3$ are closely related to the $SU(2)_k$ Kac-Moody algebra [8], [9]. Both theories can be constructed by combining a free scalar field with parafermions. In [2] it was shown that a similar relationship exists between $N = 2$ models with $c > 3$ and $SU(1, 1)$ models. In this case we combine the generalized parafermions $SU(1, 1)/U(1)$ [10] with a free scalar field. The currents satisfying the operator product expansion of the $SU(1, 1)$ Kac-Moody algebra are given by

$$\begin{aligned} J^+(z) &= \sqrt{k}\psi(z)e^{\sqrt{\frac{2}{k}}\phi}, \\ J^-(z) &= \sqrt{k}\psi^\dagger(z)e^{-\sqrt{\frac{2}{k}}\phi}, \\ J^3(z) &= -\sqrt{\frac{k}{2}}\partial_z\phi, \end{aligned} \tag{2.1}$$

where $\psi_1(z)$ and $\psi_1^\dagger(z)$ are the parafermions and the field $\phi(z)$ describes the $U(1)$ part. The supersymmetry generators are obtained by changing the coefficients of the exponential term in (2.1),

$$\begin{aligned} G^+(z) &= \sqrt{\frac{2c}{3}}\psi(z)e^{\sqrt{\frac{3}{c}}\phi}, \\ G^-(z) &= \sqrt{\frac{2c}{3}}\psi^\dagger(z)e^{-\sqrt{\frac{3}{c}}\phi}, \\ J(z) &= i\sqrt{\frac{c}{3}}\partial_z\phi. \end{aligned} \tag{2.2}$$

Using the parafermion algebra [10] one can verify that the fields G^+ , G^- , J of (2.2), along with the stress tensor $T(z) = T_\psi - \frac{1}{2}(\partial_z\phi)^2$, satisfy the $N = 2$ algebra. The relationship between the central charge c of the Virasoro algebra and the level k Kac-Moody algebra is $c = 3k/(k - 2) > 3$.

With this connection between $SU(1, 1)$ model and generalized parafermionic theory, it is clear that every field in the theory can be written as a product of a field in the parafermionic and a field in the $U(1)$ theory. The field in the corresponding $N = 2$ theory is found by changing the radius of the boson [2]. Thus, the characters of $N = 2$ superconformal unitary models with $c > 3$ can be calculated by first finding the character for the unitary $SU(1, 1)/U(1)$ model and then combining this parafermionic character with a free boson character to form a character of the unitary $N = 2$ representation. The characters for the parafermionic coset model $SU(1, 1)/U(1)$ has been calculated in [11]. When one adds back the boson to complete the $N = 2$ representation it is important to

notice that the $SU(1,1)$ character does not truncate [10] as it does in the compact case of $SU(2)_k$, where a Z_k symmetry is present.

Alternatively one can find the characters for representations of $N = 2$ superconformal highest weight representations for $c > 3$ and view them as $SU(1,1)$ representations. The characters from null state structure of unitary $N = 2$ representations have been derived in [12], [13], [14]. We use the results in [15], where the $N = 2$ characters in NS sector were interpreted as non-compact characters. The character for a highest weight $N = 2$ representation R is defined as

$$\chi_R(\tau, z, u = 0) = \text{Tr}_R[q^{L_0} w^{J_0}], \quad (2.3)$$

where $q = e^{2\pi i \tau}$ and $w = e^{2\pi i z}$. Later we specialize to the characters $\chi_R(\tau, 0, 0)$.

The dimension $h_{j,m}$ and the $U(1)$ charge $Q_{j,m}$ of an $N = 2$ primary state are given by

$$h_{j,m} = \frac{-j(j-1) + m^2}{(k-2)}, \quad Q_{j,m} = -\frac{2m}{k-2}. \quad (2.4)$$

Corresponding to $SU(1,1)$ representations [16] there are four possibilities (NS sector):

1. Trivial representation for which $j = 0$, $m = 0$ and

$$\chi^{triv}(\tau, z) = F(q, w) \frac{1 - q}{(1 + q^{1/2}w)(1 + q^{1/2}w^{-1})}, \quad (2.5)$$

where $F(q, w) = \prod_{n=1}^{\infty} (1 - q^n)^{-3} \sum_{n \in \mathbf{Z}} q^{n^2/2} w^n$.

2. Positive discrete series D_n^+ , for which $j = n + \epsilon$, $\epsilon = 0, 1/2$ and $m = j + r$, $r \in \mathbf{Z}_0^+$ and

$$\begin{aligned} j \neq k/2: \quad \chi_{j,m}^+(\tau, z) &= q^{h_{j,m}} w^{Q_{j,m}} \frac{F(q, w)}{1 + q^{r+1/2}w^{-1}}, \\ j = k/2: \quad \chi_{k/2,m}^+(\tau, z) &= q^{h_{k/2,m}} w^{Q_{k/2,m}} \frac{F(q, w)(1 - q)}{(1 + q^{r+1/2}w^{-1})(1 + q^{r+3/2}w^{-1})}. \end{aligned} \quad (2.6)$$

3. Negative discrete series D_n^- , for which $j = n - \epsilon$, $\epsilon = 0, 1/2$ and $m = -j - r$, $r \in \mathbf{Z}_0^+$ and

$$\begin{aligned} j \neq k/2: \quad \chi_{j,m}^-(\tau, z) &= q^{h_{j,m}} w^{Q_{j,m}} \frac{F(q, w)}{1 + q^{r+1/2}w}, \\ j = k/2: \quad \chi_{k/2,m}^-(\tau, z) &= q^{h_{k/2,m}} w^{Q_{k/2,m}} \frac{F(q, w)(1 - q)}{(1 + q^{r+1/2}w)(1 + q^{r+3/2}w)}. \end{aligned} \quad (2.7)$$

4. Continuous series, for which $j(j-1) < \epsilon(\epsilon-1)$. In the principal series $j = 1/2 + i\rho$, $\rho \in \mathbf{R}$ and $m = m_0 + n$, $n \in \mathbf{Z}$, $|m_0| \leq 1/2$. In the supplementary series j is real with $0 < |j - 1/2| < \epsilon + 1/2$ and $m = m_0 + n$, $n \in \mathbf{Z}$, $|m_0 \pm 1/2| < |j - 1/2|$ and

$$\chi_{j,m}^c(\tau, z) = q^{h_{j,m}} w^{Q_{j,m}} F(q, w). \quad (2.8)$$

We are interested in finding a modular invariant partition function

$$Z = \sum_{p,q} N_{p,q} \chi_p(\tau, 0, 0) \chi_q(\tau, 0, 0)^*. \quad (2.9)$$

We also require that the coefficient of the vacuum contribution in eq. (2.9) is one and the coefficients $N_{p,q}$ are non-negative integers. Note that in the partition function the characters are specialized to values $z = u = 0$. In the following we use these specialized characters. It is also useful to define the supersymmetric partition $p(\ell)$ as the multiplicity of states that are generated by operating with the modes of $N = 2$ generators on the vacuum

$$F(q, 0) = \frac{\theta_3}{\prod(1 - q^n)^3} = \sum p(\ell) q^\ell = 1 + \sqrt{q} + 3q + 6q^{3/2} + 11q^2 + \dots \quad (2.10)$$

The simple structure of the characters (2.5)-(2.8) makes it easy to express q^n , $n \in \mathbf{Z}$ as a sum of the characters. The relation is found by first noticing

$$1 = \chi^{triv}(\tau) + q - \sum_{t=1/2}^{\infty} \left(\sum_{s=0}^{2t} (s+1)(-1)^s p(t - s/2) \right) (q^t - q^{t+1}), \quad (2.11)$$

where $t \in \mathbf{Z}/2$ and $s \in \mathbf{Z}$ and the powers of q on the right hand side of eq. (2.11) are given by

$$q^{\ell' + h_{j,m}} = \chi_{\ell' + h_{j,m}}^c(\tau) + \sum_{l=\ell'+1/2}^{\infty} (-1)^n \left(\prod_{i \geq 1}^n p(\ell_i) \right) \chi_{\ell + h_{j,m}}^c(\tau). \quad (2.12)$$

Here n is defined through $\sum_{i=1}^n \ell_i = \ell - \ell'$ and $\ell', \ell, \ell_i \in \mathbf{Z}_+/2$. The possible values of j , m , and $h_{j,m}$ are given in eqs. (2.4)-(2.8). The q expansion for discrete series gives

$$q^{h_{j,m}} = \chi_{h_{j,m}}^{\pm} - \sum_{t=1/2}^{\infty} \left(\sum_{0 \leq s \leq t/(r+1/2)} p(t - s(r+1/2)) (-1)^s \right) q^{t+h_{j,m}}, \quad j \neq k/2, \quad (2.13)$$

where the powers of q on the right hand side are given either by eq. (2.12) or recursively by eq. (2.13). Using eqs. (2.11) - (2.13) enables us to express the character of a Kazama-Suzuki model with known modular properties as a sum of non-compact characters.

3. $SU(3)/(SU(2) \times U(1))$ Characters

The simplest $N = 2$ superconformal model with central charge greater than three is the Kazama-Suzuki model $SU(3)/(SU(2) \times U(1))$ with $c = 6k/(k+3)$ and $k > 3$. The character for these models has been found in [17]. In NS sector we have

$$\begin{aligned} \chi_{\lambda}^{\Lambda}(\tau, z) = & \left(\frac{\theta_3(\tau, z)}{\prod (1 - q^n)^3} \right)^2 \sum_{\sigma \in W} \epsilon(\sigma) \sum_{\sigma(t)} q^{\frac{1}{k+3}((\Lambda+t+\rho)^2 - \rho^2 - ((\lambda+\hat{\rho})^2 - \hat{\rho}^2))} w^Q \\ & \times \prod_{i=1,2} \frac{1}{1 + q^{2(\sigma(\Lambda+\rho+t) - (\lambda+\hat{\rho})) \cdot \mu_i} w^{2(\alpha_1 - \alpha_2) \cdot \mu_i}}. \end{aligned} \quad (3.1)$$

where Λ (λ) is the highest weight and ρ ($\hat{\rho}$) is half of the sum of positive roots of $SU(3)$ ($SU(2) \times U(1)$), respectively. The factor $\epsilon(\sigma)$ is -1 for odd elements of the Weyl group W and $+1$ for even elements. The subgroup $SU(2)$ has been chosen to lie in the $\alpha_1 + \alpha_2$ direction, where α_1 and α_2 are the simple roots of $SU(3)$. In this case the μ_i are the fundamental weights of $SU(3)$. The characters with the other boundary conditions for fermions can be obtained by shifting z by $1/2$, $\tau/2$ or by $\tau/2 + 1/2$.

To make a connection between $SU(3)/(SU(2) \times U(1))$ model and $SU(1,1)$ theory, we need to expand (3.1) in powers of q . Consider first the highest weight $\Lambda = (0, 0)$ of $SU(3)$ and the highest weight $\lambda = (0, 0)$ of $SU(2) \times U(1)$. In appendix A we derive the power series expansion of the character $\chi_{(0,0)}^{(0,0)}(\tau)$,

$$\begin{aligned} \chi_{(0,0)}^{(0,0)}(\tau) = & \sum_{\sigma \in W} \epsilon(\sigma) \sum_{\text{quadrants}} \epsilon_q \{ (\sum_{\substack{\ell \geq \delta_{\sigma,q} \\ \ell \in \mathbf{Z}/2}} \sum_{\substack{\ell' \geq \xi_{\sigma,q} \\ \ell' \in \mathbf{Z}/2}} + \sum_{\substack{\ell \geq \delta'_{\sigma,q} \\ \ell \in \mathbf{Z}/2 + \frac{3(k+3)\mathbf{Z}_+^2}{4}}} \sum_{\substack{\ell' \geq \xi'_{\sigma,q} \\ \ell' \in \mathbf{Z}/2 + \frac{(k+3)\mathbf{Z}_+^2}{4}}}) \hat{g}_1(\ell, \ell') \\ & + \sum_{\substack{\ell \geq \alpha_{\sigma,q} \\ \ell \in \mathbf{Z}/2}} \sum_{\substack{\ell' \geq \beta_{\sigma,q} \\ \ell' \in \mathbf{Z}/2}} \hat{g}_2(\ell, \ell') \} q^{\ell+\ell'}. \end{aligned} \quad (3.2)$$

As an example, take the $SU(3)$ coset model at level 4 in terms of the $SU(1,1)$ characters at level 16. We find $\chi_{(0,0)}^{(0,0)}(\tau)$ in terms of $SU(1,1)$ characters by using (3.2) together with (2.11) and (2.12). The explicit multiplicities of the first few terms are

$$\begin{aligned} \chi_{\lambda=(0,0)}^{\Lambda=(0,0)}(\tau, z) = & \chi^{triv} + \chi_{2,0}^c + \chi_{4,0}^c + 2\chi_{11/2,0}^c + \chi_{6,0}^c + \chi_{7,0}^c + 2\chi_{15/2,0}^c + 3\chi_{8,0}^c + \chi_{9,0}^c \\ & + 2\chi_{19/2,0}^c + 5\chi_{10,0}^c + 2\chi_{21/2,0}^c + \chi_{11,0}^c + 6\chi_{23/2,0}^c + 4\chi_{12,0}^c + 2\chi_{25/2,0}^c + \dots \end{aligned} \quad (3.3)$$

The subscripts of the continuous characters correspond to (h, Q) . We have checked with Mathematica that also the J_0 parts on the left and right hand side of eq. (3.3) are equal. This is expected, since if one makes a twist, which preserves the algebra [18],

$$\begin{aligned} L_n &\rightarrow L_n + \eta J_n + \frac{1}{6}\eta^2 c \delta_{n,0}, \\ J_n &\rightarrow J_n + \frac{1}{3}\eta c \delta_{n,0}, \\ G_r^\pm &\rightarrow G_{r\pm\eta}^\pm, \end{aligned} \tag{3.4}$$

to the $N = 2$ character $\chi(\tau, 0, 0)$, and after that a modular transformation $(\tau, z, u) \rightarrow (-1/\tau, z/\tau, u + (cz^2)/(6\tau))$ [19], one obtains the following relation between the twisted and non-twisted characters:

$$\chi(-1/\tau, \eta, 0) = \chi^\eta(\tau, 0, 0)|_S. \tag{3.5}$$

This implies that also the $U(1)$ parts in eq. (3.3) are equivalent, when the characters are evaluated at the same point.

It is also possible to deconstruct the characters of other representations explicitly similarly than (3.2). This deconstruction is given in appendix A.

4. Modular Invariants and Discussion

We have demonstrated that $SU(3)/(SU(2) \times U(1))$ characters can be written as a sum of $SU(1,1)$ characters. Since modular transformation matrices are unitary, it is well known that combining the $SU(3)/(SU(2) \times U(1))$ left and right sectors diagonally leads to a modular invariant partition function

$$Z(\tau) = \frac{1}{2} \sum |\chi_{\lambda,\nu}^\Lambda(\tau)|^2, \tag{4.1}$$

where ν denotes the different boundary conditions for the fermions. This means that also the infinite combination of specialized $SU(1,1)$ characters, given by the decomposition of χ_λ^Λ , is modular invariant.

The infinite number of $SU(1,1)_{16}$ characters in eq. (3.3) forming the character of the $k = 4$ Kazama-Suzuki model has a natural explanation. In [2] it was shown that an infinite number of representations is needed to construct a modular invariant partition function for $N = 2$ theory, if $c > 3$. This generalizes Cardy's result [20] for Virasoro minimal models. In the case of the $SU(3)/(SU(2) \times U(1))$ model, when the $q^{1/24}$ factors

of η functions and vacuum contribution $q^{-c/24}$ are properly included in the character (3.1), the maximum growth of the partition function in the limit $\tau \rightarrow i\infty$ is given by the vacuum energy $e^{\pi c \text{Im}\tau/6}$. On the other hand, after the modular transformation $\tau \rightarrow -1/\tau$ and the transformation to $\tau \rightarrow i\infty$ region, the maximum growth is $e^{\pi \text{Im}\tau}$. This shows that for $c < 6$, the Kazama-Suzuki partition function doesn't necessarily have an infinite number of terms, when $k = 4$, while for the $SU(1,1)$ model at $k = 16$ an infinite number of representations is needed.

Because of the GSO projection, the partition function for ten dimensional Neveu-Schwarz-Ramond model is known to vanish by the Riemann identity. In [21] it was shown that there exists a generalized Riemann identity by which the diagonal four dimensional partition function, based on compactification on nine copies of superconformal minimal models at level 1 vanishes. Along similar lines one should be able to explicitly show that the four dimensional partition function for Kazama-Suzuki models $SU(n)/(SU(n-1) \times U(1))$ with $c = 9$ vanishes. We hope to return to this issue later.

Acknowledgement:

I would like to thank Dennis Nemeschansky for suggestions and discussions.

Appendix A.

In this appendix we write the character as the power series expansion (3.2). The character (3.1) can be written as

$$q^{\frac{1}{p}(\Lambda+\rho)^2} \sum_{\ell, \ell'=0}^{\infty} p(\ell)p(\ell')q^{\ell+\ell'} \sum_{\sigma} \epsilon(\sigma) \left(\prod_{i=1,2} \left(\sum_{\substack{\ell_i \succ 0 \\ s_i \geq 0}} - \sum_{\substack{\ell_i \prec 0 \\ s_i < 0}} \right) \right) (-1)^{s_1+s_2} \quad (\text{A.1})$$

$$\times q^{p(\ell_1^2+\ell_2^2-\ell_1\ell_2)+2\bar{\ell}\cdot\sigma(\rho)+p\bar{\ell}\cdot\bar{s}+[\sigma(\Lambda+\rho)-\lambda-\hat{\rho}]\cdot\bar{s}}.$$

Here $\ell_i \succ 0$ means $\ell_i \geq 0$, if $(\sigma(\Lambda + \rho) - \lambda - \hat{\rho}) \cdot \mu_i > 0$, otherwise $\ell_i < 0$. If $\ell_i \geq 0$ in the first term, then $\ell_i < 0$ in the second term, otherwise $\ell_i \leq 0$. For our purpose it is useful to diagonalize the exponent in ℓ_1 and ℓ_2 . To do that we write $\ell_1 = \ell'_1 + \ell'_2$ and $\ell_2 = \ell'_1 - \ell'_2$ and $\ell'_1 \pm \ell'_2 \in \mathbb{Z}$.

With this change and changing the summation appropriately each term in the character (A.1) can be written in the form:

$$\begin{aligned} & \pm q^{\frac{1}{6p}(3a^2+b^2)} \sum_{\substack{\ell, \ell' \\ \ell, \ell' \in \mathbb{Z}/2}} p(\ell)p(\ell')q^{\ell+\ell'} \\ & \times \sum_{\substack{\ell'_1 \succ 0 \\ -\ell'_1 \prec \ell'_2 \prec \ell'_1 \\ s_1, s_2 \succ 0}} (-1)^{s_1+s_2} q^{(p(\ell'_1{}^2+3\ell'_2{}^2)+a\ell'_1+b\ell'_2)+s_1(p(\ell'_1+\ell'_2)+c)+s_2(p(\ell'_1-\ell'_2)+d)}. \end{aligned} \quad (\text{A.2})$$

We have divided ℓ'_1 and ℓ'_2 sums into four quadrants according to the sign of ℓ_1 and ℓ_2 . In $-\ell'_1 \prec \ell'_2 \prec \ell'_1$ the ' $\prec$ ' means either $\leq$ or $<$: $\ell'_1 \geq -\ell'_2$ if $c > 0$ and $\ell'_2 \leq \ell'_1$ if $d > 0$. Similarly $\ell'_1 \succ 0$ is $\ell'_1 \geq 0$ if both $c > 0$ and $d > 0$, otherwise $\ell'_1 > 0$.

We consider first (A.1) with $\Lambda = (0, 0)$ and $\lambda = (0, 0)$. The coefficients a, b, c and d for different Weyl-group elements are given in table for the first quadrant, i.e. when $\ell_1 \succ 0$ and $\ell_2 \succ 0$. In the first quadrant $s_1 \geq 0, s_2 \geq 0$.

	σ_0	σ_1	σ_2	σ_{21}	σ_{12}	σ_{121}
a	2	1	1	-1	-1	-2
b	0	-3	3	3	-3	0
c	1/2	-1/2	1/2	-1/2	-3/2	-3/2
d	1/2	1/2	-1/2	-3/2	-1/2	-3/2

For the other quadrants (we change the $\ell_i \prec 0$ summations to $\ell_i \succ 0$):

$$\begin{aligned} 2. \quad & \ell_1 \prec 0, \ell_2 \succ 0: \quad s_1 > 0, s_2 \geq 0; \quad c \rightarrow -c \\ 3. \quad & \ell_1 \succ 0, \ell_2 \prec 0: \quad s_1 \geq 0, s_2 > 0; \quad d \rightarrow -d \\ 4. \quad & \ell_1 \prec 0, \ell_2 \prec 0: \quad s_1 > 0, s_2 > 0; \quad c \rightarrow -c, d \rightarrow -d. \end{aligned} \quad (\text{A.3})$$

We want to write this in a form from which we can read off the multiplicities of various $SU(1, 1)$ characters. Let's define $\ell + s_1[(\ell'_1 + \ell'_2)p + c] = \tilde{\ell}$. From this it is easy to see that

$$s_1 \leq \frac{\tilde{\ell}}{(\ell'_1 + \ell'_2)p + c} \quad (\text{A.4})$$

and the ℓ and s_1 sums can be changed as follows

$$\begin{aligned} & \sum_{\ell \succ 0} \sum_{s_1 \succ 0} p(\ell)(-1)^{s_1} q^{\ell+s_1[(\ell'_1+\ell'_2)p+c]} \\ & = \sum_{\substack{\tilde{\ell} \geq \alpha \\ \tilde{\ell} \in \mathbb{Z}_+/2}} \sum_{0 \prec s_1 \leq \frac{\tilde{\ell}}{(\ell'_1+\ell'_2)p+c}} (-1)^{s_1} p(\tilde{\ell} - s_1[(\ell'_1 + \ell'_2)p + c]) q^{\tilde{\ell}} \equiv p_1(\tilde{\ell}, \ell'_1, \ell'_2) q^{\tilde{\ell}}, \end{aligned} \quad (\text{A.5})$$

where

$$\alpha = \begin{cases} 0, & \text{if } s_1 \geq 0 \\ p + c, & \text{if } s_1 > 0. \end{cases} \quad (\text{A.6})$$

Similar considerations for ℓ' and s_2 summations give

$$\begin{aligned} & \sum_{\ell' \geq 0} \sum_{s_2 > 0} p(\ell) (-1)^{s_2} q^{\ell' + s_2[(\ell'_1 - \ell'_2)p + d]} \\ &= \sum_{\substack{\tilde{\ell}' \geq \beta \\ \tilde{\ell}' \in \mathbf{Z}_+/2}} \left[\sum_{0 \prec s_2 \prec \frac{\tilde{\ell}'}{(\ell'_1 - \ell'_2)p + d}} (-1)^{s_2} p(\tilde{\ell}' - s_2[(\ell'_1 - \ell'_2)p + d]) q^{\tilde{\ell}'} \right] \equiv p'_1(\tilde{\ell}', \ell'_1, \ell'_2) q^{\tilde{\ell}'}, \end{aligned} \quad (\text{A.7})$$

where

$$\beta = \begin{cases} 0, & \text{if } s_2 \geq 0 \\ p + d, & \text{if } s_2 > 0. \end{cases} \quad (\text{A.8})$$

We still want to deal with the summations over ℓ'_1 and ℓ'_2 .

Consider first the case when ℓ'_1 and ℓ'_2 are integers and $\ell'_1 \neq 0$. (Note that they are always both either integers or integers+half.) The dependence of ℓ'_2 is

$$\sum_{-\ell'_1 \prec \ell'_2 \prec \ell'_1} \sum_{\tilde{\ell} \geq \alpha} p_1(\tilde{\ell}, \ell'_1, \ell'_2) p'_1(\tilde{\ell}, \ell'_1, \ell'_2) q^{\tilde{\ell} + 3\ell'^2_2 p + b\ell'_2}. \quad (\text{A.9})$$

Changing summation over the variable $\hat{\ell}$, where $\hat{\ell}$ is defined as

$$3\ell'^2_2 p + b\ell'_2 + \tilde{\ell} \equiv \hat{\ell} \in \mathbf{Z}/2 \quad (\text{A.10})$$

we get (A.9) to the following form

$$\sum_{\substack{\hat{\ell} \geq \delta \\ \hat{\ell} \in \mathbf{Z}/2}} \sum_{\tilde{\gamma}_- \prec \ell'_2 \prec \tilde{\gamma}_+} p_1(\tilde{\ell}, \ell'_1, \ell'_2) p'_1(\tilde{\ell}, \ell'_1, \ell'_2) q^{\hat{\ell}} \equiv \sum_{\substack{\hat{\ell} \geq \delta \\ \hat{\ell} \in \mathbf{Z}/2}} g(\hat{\ell}, \tilde{\ell}', \ell'_1) q^{\hat{\ell}}. \quad (\text{A.11})$$

Here $\tilde{\gamma}_{\pm} = \{\min_{\max}\} \{\pm \tilde{\ell}'_1, -b/6p \pm \sqrt{b^2 + 12p\hat{\ell} - 12\alpha p/6p}\}$ and $\tilde{\ell}'_1 = \begin{cases} \ell'_1, & \text{if } \ell'_1 \geq \pm \ell'_2 \\ \ell'_1 - 1, & \text{otherwise} \end{cases}$ and $\delta = \alpha - b^2/12p$.

Defining $\tilde{\ell}' + \ell'^2_1 p + a\ell'_1 = \hat{\ell}'$, we get from the ℓ'_1 summation

$$\sum_{\substack{\hat{\ell}' \geq \xi \\ \hat{\ell}' \in \mathbf{Z}/2}} \sum_{\max\{\tilde{\zeta}, \zeta_-\} \prec \ell'_1 \prec \zeta_+} g(\hat{\ell}, \tilde{\ell}', \ell'_1) q^{\hat{\ell}'} \equiv \sum_{\substack{\hat{\ell} \geq \delta \\ \hat{\ell} \in \mathbf{Z}/2}} \sum_{\substack{\hat{\ell}' \geq \xi \\ \hat{\ell}' \in \mathbf{Z}/2}} \hat{g}_1(\hat{\ell}, \hat{\ell}') q^{\hat{\ell}'}. \quad (\text{A.12})$$

Here $\xi = \beta - a^2/4p$, $\zeta_{\pm} = a/2p \pm \sqrt{a^2 + 4p\hat{\ell}' - 4p\beta/2p}$ and $\tilde{\zeta} = \begin{cases} 0, & \text{if } \ell'_1 \geq 0 \\ 1, & \text{otherwise} \end{cases}$.

From the above we have that when ℓ'_1 and ℓ'_2 are integers and $\ell'_1 \neq 0$, the contribution to the character is

$$\sum_{\substack{\hat{\ell} \geq \delta \\ \hat{\ell} \in \mathbf{Z}/2}} \sum_{\substack{\hat{\ell}' \geq \xi \\ \hat{\ell}' \in \mathbf{Z}/2}} \hat{g}_1(\hat{\ell}, \hat{\ell}') q^{\hat{\ell} + \hat{\ell}'} \quad (\text{A.13})$$

If ℓ'_1 and ℓ'_2 are integers+half we get otherwise similar term, but $\hat{\ell} \geq \max(\delta, \frac{3p}{4} + \frac{b}{2}) \equiv \delta'$, $\hat{\ell} \in \mathbf{Z}/2 + \frac{3pn^2}{4}$ and $\hat{\ell} \geq \max(\xi, \frac{p}{4} + \frac{a}{2}) \equiv \xi'$, $\hat{\ell} \in \mathbf{Z}/2 + \frac{pn^2}{4}$.

If finally $\ell'_1 = 0$, it is easy to see that the contribution of this term is

$$\sum_{\substack{\hat{\ell} \geq \alpha \\ \hat{\ell} \in \mathbf{Z}/2}} \sum_{\substack{\hat{\ell}' \geq \beta \\ \hat{\ell}' \in \mathbf{Z}/2}} \sum_{\substack{0 \prec s_1 \leq \hat{\ell}/c \\ 0 \prec s_2 \leq \hat{\ell}'/d}} (-1)^{s_1 + s_2} p(\hat{\ell} - s_1 c) p(\hat{\ell}' - s_2 d) q^{\hat{\ell} + \hat{\ell}'} = \sum_{\substack{\hat{\ell} \geq \alpha \\ \hat{\ell} \in \mathbf{Z}/2}} \sum_{\substack{\hat{\ell}' \geq \beta \\ \hat{\ell}' \in \mathbf{Z}/2}} \hat{g}_2(\hat{\ell}, \hat{\ell}') q^{\hat{\ell} + \hat{\ell}'}. \quad (\text{A.14})$$

Combining the previous results gives for the character

$$\begin{aligned} & \sum_{\sigma \in W} \epsilon(\sigma) \sum_{\text{quadrants}} \epsilon_q \left\{ \left(\sum_{\substack{\hat{\ell} \geq \delta_{\sigma, q} \\ \hat{\ell} \in \mathbf{Z}/2}} \sum_{\substack{\hat{\ell}' \geq \xi_{\sigma, q} \\ \hat{\ell}' \in \mathbf{Z}/2}} + \sum_{\substack{\hat{\ell} \geq \delta'_{\sigma, q} \\ \hat{\ell} \in \mathbf{Z}/2 + \frac{3p\mathbf{Z}^2}{4}}} \sum_{\substack{\hat{\ell}' \geq \xi'_{\sigma, q} \\ \hat{\ell}' \in \mathbf{Z}/2 + \frac{p\mathbf{Z}^2}{4}}} \right) \hat{g}_1(\hat{\ell}, \hat{\ell}') + \right. \\ & \quad \left. \sum_{\substack{\hat{\ell} \geq \alpha_{\sigma, q} \\ \hat{\ell} \in \mathbf{Z}/2}} \sum_{\substack{\hat{\ell}' \geq \beta_{\sigma, q} \\ \hat{\ell}' \in \mathbf{Z}/2}} \hat{g}_2(\hat{\ell}, \hat{\ell}') \right\} q^{\hat{\ell} + \hat{\ell}'}, \end{aligned} \quad (\text{A.15})$$

where the subscript q refers to specific values in the given quadrant. $\epsilon_q = 1$ for the first and fourth quadrant, otherwise it is -1 . For general Λ , λ the summation indices $\hat{\ell}, \hat{\ell}'$ contain the fractional part from the coefficients a, b, c, d .

References

- [1] Y. Kazama, H. Suzuki, Nucl. Phys. B321 (1989) 232.
- [2] L. Dixon, M. Peskin, J. Lykken, Nucl. Phys. B 325 (1989) 329.
- [3] I. Bars, Nucl. Phys. B334 (1990) 125.
- [4] I. Bars, D. Nemeschansky, Nucl. Phys. B348 (1991) 89.
- [5] A. N. Schellekens, S. Yankielowicz, Int. J. Mod. Phys. A5 (1990) 2903.
- [6] M. Henningson, S. Hwang, P. Roberts and B. Sundborg, Phys. Lett. B 267 (1991) 350.
- [7] E. Witten, Phys. Rev. D44 (1991) 314.
- [8] A. B. Zamolodchikov, V. A. Fateev, Sov. Phys. JETP 63 (1986) 913.
- [9] Z. Qiu, Phys. Lett. 198B (1987) 497.
- [10] J. Lykken, Nucl. Phys. B313 (1989) 473.
- [11] P. Griffin, Nucl. Phys. B356 (1991) 287; K. Sfetsos, Phys. Lett. B271 (1991) 301.
- [12] V. K. Dobrev, Phys. Lett. 186 B (1987) 43.
- [13] Y. Matsuo, Prog. Theor. Phys. 77 (1987) 7930.
- [14] E. Kiritsis, Int. J. Mod. Phys. A3 (1988) 1871.
- [15] I. Bakas, E. Kiritsis, UCB-PTH-91/44.
- [16] V. Bargmann, Ann. Math. 48 (1947) 568, Commun. Pure Appl. Math. 14 (1961) 187.
- [17] K. Huitu, D. Nemeschansky, preprint USC-92/011.
- [18] A. Schwimmer, N. Seiberg, Phys. Lett. 184 B (1987) 191.
- [19] D. Gepner, E. Witten, Nucl. Phys. B 278 (1986) 493.
- [20] J. Cardy, Nucl. Phys. B270 [FS16] (1986) 186.
- [21] S. K. Han, Phys. Rev. D 39 (1989) 2322.